

Biox® -15EC Line Spray

P O S T H A R V E S T

Potato Sprout Control



ACTIVE INGREDIENTS:

Clove Oil	15.0 %
INERT INGREDIENTS: (Glycerin, Potassium oleate, water).....	85.0 %
TOTAL.....	100.0 %

NOTICE: This product has not been registered by the US Environmental Protection Agency (EPA). Pace International, LLC represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA).



KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO
Si usted no entiende la etiqueta, busque a alguien para que se la explique en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

Danger. Harmful if swallowed. Maybe harmful if inhaled. Causes skin and eye irritation. May cause respiratory irritation and allergy or asthma symptoms or breathing difficulties if inhaled. May cause damage to organs (nervous system, mucous membranes, liver, blood forming systems (bone marrow and spleen) through prolonged or repeated exposure. Do not breathe mist, vapors, or spray. In case of inadequate ventilation wear respiratory protection, best used outdoors or in a well-ventilated area. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective clothing, chemical resistant gloves, and safety glasses.

FIRST AID

IF SWALLOWED: Call a POISON CENTER / doctor if you feel unwell. Rinse mouth. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Call a POISON CENTER / doctor if you feel unwell or are experiencing respiratory symptoms. Remove person to fresh air and keep comfortable for breathing.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. If eye irritation persists: Get medical advice/attention.

IF ON SKIN: Wash with plenty of soap and water. Take off contaminated clothing and wash it before reuse. If skin irritation occurs: Get medical advice/attention. See First Aid, Section 4 on your SDS for further instructions.

NOTE: Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection sheet.

Applicators and other handlers must wear: long-sleeved shirt, long pants, shoes, socks, protective eyewear (goggles or face shield), and chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber or butyl rubber. Wash any contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

DIRECTIONS FOR USE

This product is used for treating potatoes to suppress sprouting after storage and washing. This product will not, if used in accordance with the label, prevent wound periderm formation of potatoes.

DO NOT APPLY IN THE FIELD.

IMPORTANT: Potatoes may deteriorate or otherwise be damaged by bacteria, fungi, insects, physiological aging and physical effects such as internal sprouting, environmental factors (such as humidity, temperature and airflow), and other causes which are unrelated to the application of this product, and which may occur whether or not this product is applied to the potatoes.

DO NOT USE ON SEED POTATOES: This product will kill emerging buds on potato tubers and therefore should NOT be used on potatoes that will be used for seed. The damage to the emerged buds is permanent although secondary buds may grow after 30 to 90 days. The time period of effective sprout suppression may vary depending on many conditions beyond the control of the manufacturer and/or seller, including, but not limited to, timing and method of application, variety and condition of stored crop, and mixtures or treatments with other chemicals. Wash contaminated equipment thoroughly with soap and hot water before use on seed potatoes.

EQUIPMENT: Biox®-15EC is usually applied as a 0.8% to 1.8% emulsion wash on the drying table, or similar station, after the potatoes have been washed. Equipment should be set to apply said emulsion evenly on the potatoes moving across the rollers. Rolling or tumbling action greatly improves coverage compared to flat conveyor belts. Thorough coverage of the potato by the emulsion will provide the best results. For further information about application equipment, contact your local Biox® Distributor Representative.

APPLICATION: Proper treatment solutions of 0.8% to 1.8% solutions (Column A) may be prepared by adding Column B to Column C:

Column A	Column B	Column C
*Emulsion concentration	Amt of Product (gal)	Amt of Water (gal)
0.83%	(8,300 ppm)	1 20.0
1.00%	(10,000 ppm)	1 16.5
1.21%	(12,100 ppm)	1 13.5
1.40%	(14,000 ppm)	1 11.5
1.59%	(15,900 ppm)	1 10.0
1.84%	(18,400 ppm)	1 8.5

* These percentages are based on w/vol ratio and are rounded to the nearest 100 ppm.

NOTE: Thorough agitation at initial mixing is required – no further agitation needed after the emulsion is completely mixed.

For a flood application system, use a sufficient emulsified solution of the desired concentration and apply in a method to provide thorough


For a flood application system, use a sufficient emulsified solution of the desired concentration and apply in a method to provide thorough coverage/wetting of the potatoes. The effectiveness of this product is not only based on uniform coverage but a minimum residue concentration of approximately 15 parts per million (ppm) should maintain on the potatoes after treatment. In addition, the extent of peeper/sprout development, the cleanness of the potatoes at the time the solution is applied, the potato variety (russet vs. smooth skin types) and the wetness of the tubers after washing will also affect the performance of this product. It is suggested that residue analysis be conducted on a routine bases to ensure that the desired active ingredient is achieved on the potatoes.



Manufactured by:

Pace International LLC

5661 Branch Road, Wapato, WA 98951 800.936.6750



Biox® -15EC

Line Spray

P O S T H A R V E S T

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Pesticide Storage: Biox®-15EC and the emulsified solution (once completely mixed) can be stored at room temperature. Store in a well-ventilated place. Keep container tightly closed when not in use. Store locked up. This product temporarily inhibits germination of seed potatoes.

Pesticide Disposal: Wastes resulting from the use of the product may be disposed of on site or in accordance with applicable regional, national and local laws and regulations.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

011218US

The use of Biox®-15EC in agricultural crop protection is covered by US Patent 5,858,436 and other patents pending.
Biox is a registered trademark of Pace International, LLC.
Product of USA

Specimen Label